arXiv:1102.0995v2 [astro-ph.CO] 4 May 2011

Observational constraints of the Gravitational Waves in the Brans-Dicke Theory

R. C. Freitas* and S. V. B. Gonçalves†

Universidade Federal do Espírito Santo, Centro de Ciências Exatas, Departamento de Física

Av. Fernando Ferrari, 514 - Campus de Goiabeiras, CEP 29075-910, Vitória, Espírito Santo, Brazil

January 11, 2013

Abstract

We investigate the quantum origin of the primordial cosmological gravitational waves in the Brans-Dicke theory. We compute the number of gravitons N_k produced during inflation and the observables: power spectrum P_T, spectral index n_T and energy density Ω_k. By comparison with General Relativity we see that the results for both theories are the same for the case of the particles number N_k and for the case of the power spectrum P_T and the energy density Ω only when $\omega < 10$. For the spectral index n_T we found that when the Brans-Dicke coupling parameter ω is bigger than unity the spectral index approximates the expression found in the General Relativity. This may awake us to the possibility that ω varies with the cosmological scale.

KEYWORDS: scalar-tensor gravity, gravitational waves

PACS numbers: 04.30.-w, 98.80.-K

1 Introduction

The gravitational waves are tensorial fluctuations in the metric of spacetime. This particular perturbation is not explicitly coupled with the energy density and pressure of the matter of the Universe and does not contribute to the gravitational instability that generates the cosmological structures we see today. On the other side his study is of great interest because it supplies the specific signature of the metric theory of gravity. Nevertheless, they must have left special signatures in the polarization of the CMB anisotropies [1–3]. Moreover, since the gravitational waves were generated in the early Universe, before the time of last scattering of CMB photons, they can be a window to the primordial phase of evolution of the Universe. These waves are

*e-mail: rc_freitas@terra.com.br

†e-mail: sergio.vitorino@pq.cnpq.br

predicted by Einstein's theory of General Relativity, but they still have to be directly detected. Great efforts are been done in this sense and there is hope that a new generation of experiments in space may allow this detection in some years [4]. The gravitational waves spectrum frequency extends over a wide range of interest, from $10^{-18}Hz$ to $10^{8}Hz$[1], depending of the sources that generate those waves. Many works have been made in order to identify specific signatures of cosmological models in the spectra of gravitational waves, for example, in the case of quintessence model [5] and string cosmology [6–8].

The existence of a classical scalar field in the nature has been considered in many theories of gravitation that present alternatives to General Relativity. The prototype of scalar-tensor theories is the Brans-Dicke theory [9–12], whose Lagrangian in the Jordan frame is given by

$$\mathscr{S} = \frac{1}{16\pi}\int d^4x\sqrt{-g}\left[\phi R - \omega\left(\frac{\phi_{,\mu}\phi^{,\mu}}{\phi}\right) + 16\pi\mathscr{L}_{mat}\right] \quad , \tag{1}$$

where ϕ is a scalar field that couples to gravity through the parameter ω, called Brans-Dicke parameter and $\mathscr{L}_{mat}$ is the matter Lagrangian.

The Brans-Dicke theory, as others scalar-tensor theories, can be formulated in the Einstein or in the Jordan frames, which are conformally related. Which conformal frame is the physical one is a very contentious issue [13]. The most common argument against the Jordan frame found in the literature is that the scalar field energy density can assume negative values. This happens because of the nonmininal coupling between the scalar field and geometry. The terms with the second covariant derivatives of the scalar field in the Brans-Dicke field equations contain the connection $\Gamma^{\alpha}_{\mu\nu}$ and therefore part of the dynamical description of gravity. But the energy density can be made nonnegative with the help of a new connection $\tilde{\Gamma}^{\alpha}_{\mu\nu}$ [14] which is also associated with the physical metric $g_{\mu\nu}$. Based on the considerations presented in the previous reference we work here considering Jordan frame as the physical frame.

One can expect that the presence of the scalar field leads to different predictions about the results obtained in General Relativity. The standard cosmological scenario given by General Relativity shows some spectacular successes but also have some puzzles, such as the flatness problem, the horizon problem and structure formation problems [15].

The value of the parameter ω can be limited by local physics experiments like Cassini experiment [16, 17]: it should be huge today, the order of $\omega = 40000$ to satisfy the experimental tests of General Relativity. This result reduced the interest in the Brans-Dicke theory because when $\omega \to \infty$ in the motion equations we obtain the General Relativity. The situation has changed with the proposal of the extended inflation [18, 19]. The idea is to take advantage of the time dependence of gravitational constant G to solve the bubble nucleation problem that arises in tradicional inflation. In the de Sitter phase, Brans-Dicke theory predicts power-law inflation instead of exponential; in fact, following [18], at the beginning of inflation the Brans-Dicke solutions (for $p = -\rho$) approaches the Einstein-de Sitter solution. In the second stage of the inflation, both the scalar field and the scale factor grow by power law rather than exponential. This feature prevents the so-called graceful exit problem. However, in order to work, the parameter must be $\omega \approx 24$, contradicting local observations. This drawback can be overcome through a generalization of the original Brans-Dicke theory, allowing the parameter ω to be a function of the field ϕ itself.

[1]The wave number interval is from $10^{-26}m^{-1}$ to $10^{-16}m^{-1}$.

In 1999, the SN Ia observations [20] showed that the universe is currently undergoing accelerated expansion. A possible theoretical explanation for this acceleration is the vaccum energy with negative pressure, called dark energy (that violates the strong energy condition). There are many dark energy models, for example quintessence [21], k-essence [22], phantom [23] and Chaplygin gas [24]. The simplest candidate for the dark energy is the cosmological constant Λ whose equation of state is $p = -\rho$. But there are some problems with this alternative: why the cosmological constant is so small, nonzero, and comparable to the critical density at the present? So, it is interesting to study dynamical dark energy models like the Brans-Dicke theory of gravity as a possible theory of k-essence coupled to gravity since it involves the simplest form of non-linear kinetic term for the Brans-Dicke scalar field $\phi(t)$ [25].

This revival of the Brans-Dicke theory leads us to ask if the generation of gravity waves can be modified through the introduction of this scalar field. Here, we propose to study the evolution of tensorial fluctuation in the traditional Brans-Dicke theory and its quantum origin and calculate the observables that can be compared with the observational data. The main interest in the presence of the scalar field is the possibility that it can change the evolution of gravity waves in comparison with the results obtained with General Relativity [26].

This paper is organized as follows. In section II we show the field equations, the equations of motion of the Brans-Dicke theory and the background solutions. In section III we obtain the perturbed equations of gravity waves and their solutions in terms of Hankel functions in the primordial phase of the universe. In section IV the quantum formulation of the gravitational waves in the Brans-Dicke theory is obtained. In section V we calculate the observables of the Brans-Dicke theory. Finally, in section VI we discuss the results and show our conclusions.

We use in this article standard tensor notation: the greek indices run from zero to three; the latin indices run from one to three; the space-time metric is taken to have signature $+2$, or $(-,+,+,+)$; covariant derivatives are indicated by semicolons; partial derivatives are indicated by commas and the scalar field ϕ is a time function.

2 Background solutions

We assume the background Universe is spatially flat, homogeneous and isotropic, *i.e.*, it is described by the Robertson-Walker metric

$$\mathrm{d}s^2 = -\mathrm{d}t^2 + a^2(t)\left(\mathrm{d}x^2 + \mathrm{d}y^2 + \mathrm{d}z^2\right) \ , \tag{2}$$

where the $a(t)$ is the scale factor of the Universe, and $c = 1$. The energy-momentum tensor of the background matter takes a perfect fluid form

$$T^{\mu\nu} = (\rho + p)\, u^\mu u^\nu + p g^{\mu\nu} \ , \tag{3}$$

with an equation of state $p = \alpha\rho$. In these expressions ρ is the matter density, p is the pressure and α is a constant. From the action (1) we obtain the field equations

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \frac{8\pi}{\phi} T_{\mu\nu} + \frac{\omega}{\phi^2}\left(\phi_{;\mu}\phi_{;\nu} - \frac{1}{2} g_{\mu\nu}\phi_{;\rho}\phi^{;\rho}\right) + \frac{1}{\phi}\left(\phi_{;\mu;\nu} - g_{\mu\nu}\Box\phi\right) \ , \tag{4}$$

$$\Box\phi = \frac{8\pi}{3+2\omega} T \ . \tag{5}$$

Inserting the metric (2), into equations (4) and (5) we obtain the equations of motion

$$-3\frac{\ddot{a}}{a} = \frac{8\pi}{\phi}\rho\left(\frac{2+\omega+3\alpha+3\alpha\omega}{3+2\omega}\right)+\omega\frac{\dot{\phi}^2}{\phi^2}+\frac{\ddot{\phi}}{\phi} \quad , \tag{6}$$

$$\ddot{\phi}+3\frac{\dot{a}}{a}\dot{\phi} = \frac{8\pi}{3+2\omega}\rho(1-3\alpha) \quad . \tag{7}$$

The above equations must be supplemented by a conservation equation of the energy-momentum tensor

$$T^{\mu\nu}_{;\nu} = 0 \quad . \tag{8}$$

When $\mu = 0$ we obtain the energy conservation

$$\dot{\rho}+3\frac{\dot{a}}{a}\rho(1+\alpha) = 0 \quad , \tag{9}$$

while, for $\mu = j$ we get the condition

$$\rho_{,i} = 0 \quad . \tag{10}$$

Background solutions [27] can be obtained assuming that the scale factor $a(t)$ and the scalar field $\phi(t)$ have a power-law form

$$a(t) \propto t^r \quad , \quad \phi(t) \propto t^s \quad . \tag{11}$$

By direct substitution of the above relation in the equations (6), (7), (9) and (10), we have

$$-3r(r-1) = 8\pi\rho t^{2-s}\left(\frac{2+\omega(1+3\alpha)+3\alpha}{3+2\omega}\right)+s^2(1+\omega)-s \quad , \tag{12}$$

$$s^2+s(3r-1) = \frac{8\pi}{3+2\omega}\rho t^{2-s}(1-3\alpha) \quad , \tag{13}$$

$$\dot{\rho}+\ 3\frac{r}{t}\rho(1+\alpha) = 0 \quad , \tag{14}$$

$$\rho_{,i} = 0 \quad . \tag{15}$$

In what follows we shall consider the inflation case ($\alpha = -1$), the radiation phase ($\alpha = 1/3$) and the dust era ($\alpha = 0$). In the first scenario we have a drastic expansion of the Universe during the early period of the Big Bang. In this case, the background solutions are

$$s = 2 \ , \quad r = \omega+\frac{1}{2} \quad . \tag{16}$$

In the radiation phase, with $\alpha = 1/3$, we have the same background solution that the General Relativity

$$s = 0 \ , \quad r = \frac{1}{2} \quad . \tag{17}$$

In the dust era ($\alpha = 0$) we obtain

$$s = 2-3r \ , \quad r = 2\frac{1+\omega}{4+3\omega} \quad , \tag{18}$$

3 The gravitational waves equation in the Brans-Dicke theory

Cosmological gravitational waves are obtained by means of a small correction h_{ij} in equation (2), which represents the metric. Hence, the general expression of the metric (2), related to the unperturbed metric, is replaced by

$$\mathrm{d}s^2 = -\mathrm{d}t^2 + \left[a^2(t)\delta_{ij} + h_{ij}\right]\mathrm{d}x^i\mathrm{d}x^j \ . \tag{19}$$

Appling the metric (19) in the field equation (4) the resulting expression is

$$\ddot{h}_{ij} + \left(\frac{\dot{\phi}}{\phi} - \frac{\dot{a}}{a}\right)\dot{h}_{ij} + \left[\frac{k^2}{a^2} - 2\frac{\ddot{a}}{a} - 2\frac{\dot{a}\dot{\phi}}{a\phi}\right]h_{ij} = 0 \ , \tag{20}$$

where k is the wavenumber, the dots indicate cosmic time derivatives and we have written $h_{ij}(t,\vec{x}) = h(t)Q_{ij}$, where Q_{ij} are the eigenmodes of the Laplacian operator, such that $Q_{ii} = Q_{ki,k} = 0$.

Performing the transformation from the cosmic time to the conformal time, $a(t)\ d\eta = dt$, and representing the derivatives with respect to η by primes, equation (20) assumes the form

$$h''_{ij} + \left(\frac{\phi'}{\phi} - 2\frac{a'}{a}\right)h'_{ij} + \left[k^2 - 2\frac{a''}{a} + 2\frac{a'^2}{a^2} - 2\frac{a'\phi'}{a\phi}\right]h_{ij} = 0 \ . \tag{21}$$

To resolve this differential equation we need to perform the Fourier transformation

$$h_{ij}(\vec{x},\eta) = \sqrt{16\pi}\sum_{\lambda=\otimes,\oplus}\int \frac{\mathrm{d}^3k}{(2\pi)^{3/2}}\varepsilon_{ij}^{(\lambda)}(\hat{k})\frac{a(\eta)\mu_{(\lambda)}(\eta)}{\sqrt{\phi(\eta)}}e^{-i\vec{k}\cdot\vec{x}} \ , \tag{22}$$

where the polarization tensor $\varepsilon_{ij}^{(\lambda)}(\hat{k})$ can be decomposed as $\varepsilon_{ij}^{(\lambda)}(\hat{k})\varepsilon^{ij(\lambda')}(\hat{p}) = 2\delta^{\lambda\lambda'}\delta^{(3)}(\vec{k} - \vec{p})$ to the two polarization states $\otimes$ e $\oplus$. Then, we have

$$\mu''_{(\lambda)}(\eta) + \left[k^2 + \frac{a''}{a} + \frac{1}{2}\frac{\phi''}{\phi} - \frac{1}{4}\left(\frac{\phi'}{\phi}\right)^2 + \frac{a'\phi'}{a\phi}\right]\mu_{(\lambda)}(\eta) = 0 \ . \tag{23}$$

In this way, we rewrote the equation (21) in terms of a simple harmonic oscillator.

We can express the equation (23) in terms of the new parameter $\Phi(\eta) \equiv a(\eta)\sqrt{\phi(\eta)}$. Hence, the equation (23) becomes

$$\mu''_{(\lambda)}(\eta) + \left[k^2 - \frac{\Phi''(\eta)}{\Phi(\eta)}\right]\mu_{(\lambda)}(\eta) = 0 \ . \tag{24}$$

With the background solutions of the scale factor and the scalar field we get

$$\frac{\Phi''(\eta)}{\Phi(\eta)} = \begin{cases} \frac{F(\omega)}{|\eta|^2} \ , & \text{if} \quad \alpha = -1 \quad \text{with} \quad -\infty < \eta \leq -\eta_1 \quad , \\ 0 \ , & \text{if} \quad \alpha = 1/3 \quad \text{with} \quad -\eta_1 \leq \eta \leq \eta_2 \quad , \\ \frac{G(\omega)}{(\eta - \tilde{\eta}_{dust})^2} \ , & \text{if} \quad \alpha = 0 \quad \text{with} \quad \eta \geq \eta_2 \quad , \end{cases}$$

where

$$F(\omega) = \frac{2(3+2\omega)(1+2\omega)}{(1-2\omega)^2} \quad ,$$
$$G(\omega) = \frac{(3+2\omega)(1+\omega)}{(2+\omega)^2} \quad , \tag{25}$$

and

$$\tilde{\eta}_{dust} = \eta_2 - \frac{2+2\omega}{2+\omega}\left[1+\left(\frac{\omega+1/2}{\omega-1/2}\right)\left(1+\frac{\eta_2}{\eta_1}\right)\right] \quad , \tag{26}$$

The solutions, for each stage of evolution of the Universe are

1. Inflation era ($\alpha = -1$)

$$\mu(k\eta) = \sqrt{\eta}\left(A_1\mathscr{H}_\nu^{(1)}(k|\eta|) + A_2\mathscr{H}_\nu^{(2)}(k|\eta|)\right) \, , \tag{27}$$

where A_1 and A_2 are integration constants, $\mathscr{H}^{(1)}$ and $\mathscr{H}^{(2)}$ are the Hankel functions of first and second kind, respectively, and $\nu = \sqrt{F(\omega)+1/4}$ is the order of the Hankel functions.

2. Radiation era ($\alpha = 1/3$)

$$\mu(k\eta) = B_1 e^{ik\eta} + B_2 e^{-ik\eta} \quad , \tag{28}$$

where B_1 and B_2 are yet integration constants.

3. Dust era ($\alpha = 0$)

$$\mu(k\eta) = \sqrt{\eta}\left[C_1\mathscr{H}_\alpha^{(1)}\left(k(\eta-\tilde{\eta}_{dust})\right) + C_2\mathscr{H}_\alpha^{(2)}\left(k(\eta-\tilde{\eta}_{dust})\right)\right] \, , \tag{29}$$

where C_1 and C_2 are integration constants and $\alpha = \sqrt{G(\omega)+1/4}$ is the order of the Hankel functions.

4 The quantum gravitational waves equation in the Brans-Dicke theory

In this process we start with the Lagrangian density of the gravitational waves in the Brans-Dicke theory, given by

$$\mathscr{L} = \frac{\phi}{16\pi}\bar{g}^{\mu\nu}\partial_\mu h_{ij}\partial_\nu h^{ij} \, , \tag{30}$$

that leads to two corresponding Hamiltonians

$$H^{(1)} = -\frac{1}{2}\int \mathrm{d}^3x\left[\tilde{\pi}^2 - 2\frac{\Phi'}{\Phi}\mu\tilde{\pi} + \delta^{ij}\mu_{,i}\mu_{,j}\right] \quad , \tag{31}$$

$$H^{(2)} = -\frac{1}{2}\int \mathrm{d}^3x\left[\pi^2 - \frac{\Phi''}{\Phi}\mu^2 + \delta^{ij}\mu_{,i}\mu_{,j}\right] \, , \tag{32}$$

where π and $\tilde{\pi}$ are the canonical momentum. The above Hamiltonians differ each other by addiction or subtraction of a total conformal time derivative.

The expression of the quantum Hamiltonian obtained by the equation (32) for the gravitational waves in the Brans-Dicke theory is constructed by using the canonical commutation relations

$$[\hat{\mu}(\vec{x},\eta),\hat{\pi}(\vec{y},\eta)] = i\delta^{(3)}(\vec{x}-\vec{y}) \ . \tag{33}$$

So, by transforming the normal modes of oscillation of the gravitational waves in field and momentum operators in the Heisenberg description of quantum mechanics we obtain the Hamiltonian

$$H^{(2)} \to \hat{H}^{(2)} = -\frac{1}{2}\int \mathrm{d}^3x \left[\hat{\pi}^2 - \frac{\Phi''}{\Phi}\hat{\mu}^2 + \delta^{ij}\hat{\mu}_{,i}\hat{\mu}_{,j}\right] \ , \tag{34}$$

where the field operator is constructed from its classical field as follows

$$h_{ij} \to \hat{h}_{ij}(\vec{x},\eta) = \frac{\sqrt{16\pi}}{a(\eta)\sqrt{\phi(\eta)}}\sum_{\lambda=\otimes,\oplus}\frac{1}{2}\int \frac{\varepsilon_{ij}^{(\lambda)}(\vec{k})\mathrm{d}^3k}{(2\pi)^{3/2}}\left(\hat{\mu}_{\vec{k},(\lambda)}e^{-i\vec{k}\cdot\vec{x}} + \hat{\mu}^{\dagger}_{\vec{k},(\lambda)}e^{i\vec{k}\cdot\vec{x}}\right) \ , \tag{35}$$

with the operator $\hat{\mu}_{\vec{k}}$ given by

$$\hat{\mu}_{\vec{k}} = \hat{a}_{\vec{k}}(\eta_0)f_k(\eta) + \hat{a}^{\dagger}_{-\vec{k}}(\eta_0)f_k^*(\eta) \ , \tag{36}$$

and the function f_k satisfying the following equation

$$f_k'' + \left(k^2 - \frac{\Phi''}{\Phi}\right)f_k = 0 \ , \tag{37}$$

that is the same differential equation that describe the behavior of the classical function $\mu(\eta)$.

Now, to obtain coherent states of the relic gravitons in the Brans-Dicke theory we consider the creation and annihilation operators

$$\hat{a}_{\vec{k}} = \sqrt{\frac{k}{2}}\left(\hat{\mu}_{\vec{k}} + \frac{i}{k}\hat{\pi}_{\vec{k}}\right) \quad , \quad \hat{a}_{-\vec{k}} = \sqrt{\frac{k}{2}}\left(\hat{\mu}_{\vec{k}} - \frac{i}{k}\hat{\pi}_{\vec{k}}\right) \ , \tag{38}$$

whose temporal evolution is given by the Heisenberg equation and by the Hamiltonian (31), writen now as operator

$$H^{(1)} \to \hat{H}^{(1)} = -\frac{1}{2}\int \mathrm{d}^3x \left[\hat{\pi}^2 - \frac{\Phi'}{\Phi}\left(\hat{\mu}\hat{\pi} + \hat{\pi}\hat{\mu}\right) + \delta^{ij}\hat{\mu}_{,i}\hat{\mu}_{,j}\right] \ , \tag{39}$$

with the general solutions given by

$$\begin{aligned} \hat{a}_{\vec{k}}(\eta) &= u_k(\eta)\hat{a}_{\vec{k}}(\eta_0) + v_k(\eta)\hat{a}^{\dagger}_{-\vec{k}}(\eta_0) \quad , \\ \hat{a}^{\dagger}_{-\vec{k}}(\eta) &= v_k^*(\eta)\hat{a}_{\vec{k}}(\eta_0) + u_k^*(\eta)\hat{a}^{\dagger}_{-\vec{k}}(\eta_0) \quad , \end{aligned} \tag{40}$$

where η_0 is a fixed initial time and the functions $u_k(\eta)$ and $v_k(\eta)$ behave as

$$\frac{\mathrm{d}u_k}{\mathrm{d}\eta} = iku_k + \frac{\Phi'}{\Phi}v_k^* \quad , \quad \frac{\mathrm{d}v_k}{\mathrm{d}\eta} = ikv_k + \frac{\Phi'}{\Phi}u_k^* \quad . \tag{41}$$

The equations (40) are the Bogoliubov transformations [28] of the gravitational waves in the Brans-Dicke theory and $u_k(\eta)$ and $v_k(\eta)$ are the Bogoliubov coefficients that satisfy the relation

$$|u_k|^2 - |v_k|^2 = 1 \quad , \tag{42}$$

result that guarantees the unity of the temporal evolution of these operators.

The number operator is defined as $N_k = v_k^* v_k$. So, in our model the number of gravitons is given by

$$N_k = |v_k|^2 = \frac{|f_k|^2}{2k}\left(k^2 + \left(\frac{\Phi'}{\Phi}\right)^2\right) + \frac{1}{2k}\left|\frac{\mathrm{d}f_k}{\mathrm{d}\eta}\right|^2 - \frac{1}{2k}\frac{\Phi'}{\Phi}\left(f_k\frac{\mathrm{d}f_k^*}{\mathrm{d}\eta} + f_k^*\frac{\mathrm{d}f_k}{\mathrm{d}\eta}\right) - \frac{1}{2} \quad , \tag{43}$$

where the relation between v_k and f_k is

$$v_k^* = \frac{i}{\sqrt{2k}}\frac{\mathrm{d}f_k}{\mathrm{d}\eta} + \frac{f_k}{\sqrt{2k}}\left(k - i\frac{\Phi'}{\Phi}\right) \quad . \tag{44}$$

When $k\eta >> 1$ the number operator simplifies as

$$N_k \approx k|f_k|^2\left(1 + \frac{1}{2k^2}\left(\frac{\Phi'}{\Phi}\right)^2\right) - \frac{1}{2} \quad . \tag{45}$$

If $k\eta << 1$ we have

$$N_k \approx k|f_k|^2 - \frac{1}{2} \quad . \tag{46}$$

In Fig. 1 - Fig. 4 we show the behavior of the number of particles in the inflation era in the Brans-Dicke theory to different values of ω and small and big values of the wave number and we compare with the General Relativity. The aim here is, first of all, see how the Brans-Dicke model behaves for different values of ω, since local observations show that ω is big and, as we have written before, small values should be more interesting in large scales. Secondly, when gravitational waves of cosmological origin are observed, the Brans-Dicke theory and General Relativity may be confronted regarding the evolution of the tensor modes.

5 Observables in the Brans-Dicke theory

In this section we apply the formulation of quantum gravitational waves in Brans-Dicke theory to determine the most common and interesting observables in cosmology. The experimental data at present do not allow us to include or exclude the presence of a primordial spectrum of relic gravitons compatible with any model. The idea is that these theoretical results calculated here might in the near future be compared with the observational data obtained by the projects related to the search for gravitational waves of cosmological origin.




Figure 1: The number of gravitons produced by inflation in the Brans-Dicke theory for different values of ω, and also in the General Relativity, for the wavenumbers $k = 10^{-5}\ m^{-1}$ and $k = 10^{-1}\ m^{-1}$.




Figure 2: Same as the previous figure, but for wavenumbers $k = 10^6\ m^{-1}$ e $k = 10^7\ m^{-1}$.




Figure 3: The number of gravitons for small values of ω at two different comoving times.

5.1 Power Spectrum

The power spectrum is related by the Fourier transform of the two-point correlation function of the tensor modes in the following way [29]

$$\langle 0|\,\hat{h}^{\dagger}_{ij}(\vec{x},\eta)\hat{h}^{ij}(\vec{y},\eta)\,|0\rangle = \int P_T(k,\eta)\frac{\sin(kr)}{kr}\mathrm{d}(\ln k)\ , \tag{47}$$




Figure 4: The number of gravitons for large values of ω and for General Relativity.

where $P_T(k,\eta)$ is the power spectrum and the state $|0\rangle$ is annihilated by $\hat{a}_{\vec{k}}$.

The process of quantization of gravitational waves makes the classical field h_{ij} to be transformed into an operator $\hat{h}_{ij}$. This operator is given by the equation (35) where the vacuum state $|0\rangle$, in which there are no particles, is annihilated by the annihilation operator

$$\hat{a}_{\vec{k}}|0\rangle = 0 \quad . \tag{48}$$

The calculations of the expectation value of the field operators yields

$$\langle 0|\,\hat{h}^{\dagger}_{ij}(\vec{x},\eta)\hat{h}^{ij}(\vec{y},\eta)\,|0\rangle = 64\pi\int\frac{\mathrm{d}^3k}{(2\pi)^3}\frac{|f_k(\eta)|^2}{a^2\phi}e^{-i\vec{k}\cdot(\vec{x}-\vec{y})} \quad . \tag{49}$$

Knowing that gravitons can be produced in isotropic models and using spherical coordinates, we have

$$\langle 0|\,\hat{h}^{\dagger}_{ij}(\vec{x},\eta)\hat{h}^{ij}(\vec{y},\eta)\,|0\rangle = 64\pi\int\frac{k^2}{(2\pi)^2}\frac{|f_k|^2}{a^2\phi}\frac{2\sin(kr)}{kr}\mathrm{d}k \ , \tag{50}$$

Comparing the equation (47) with the previous equation we have, within the Brans-Dicke model, the power spectrum of the tensor modes parametrized as

$$P_T(k,\eta) = 32\pi\frac{k^3}{\pi^2}\frac{|f_k|^2}{a^2\phi} \quad . \tag{51}$$

In the inflation phase the function $f_k(k|\eta|)$ is defined as

$$f_k(k|\eta|) = \frac{\sqrt{\pi}}{2}\sqrt{|\eta|}\mathscr{H}^{(2)}_{\nu}(k|\eta|) \quad , \tag{52}$$

and by using the small argument limit of the Hankel functions [30], we have

$$P_T(k,\eta) = \frac{8\pi 2^{2\nu}\Gamma^2(\nu)}{\pi^3}\frac{k^2}{a^2\phi}(k|\eta|)^{1-2\nu} \quad . \tag{53}$$




Figure 5: The power spectrum of the gravitational waves in the Brans-Dicke theory as a function of wavenumber for small values of ω.




Figure 6: The power spectrum of the gravitational waves in the Brans-Dicke theory for large values of ω.

By definition $H = \frac{\dot{a}}{a} = \frac{1}{a^2}\frac{\mathrm{d}a}{\mathrm{d}\eta}$, and the conformal time in this model can be written as

$$|\eta| = \left(\frac{2\omega+1}{2\omega-1}\right)\frac{1}{aH} \quad . \tag{54}$$

Thus, the power spectrum in the Brans-Dicke theory during inflation for the modes out of the horizon becomes

$$P_T(k,\eta) = \frac{8\pi 2^{2\nu}\Gamma^2(\nu)}{\pi^3}\left(\frac{2\omega+1}{2\omega-1}\right)^{1-2\nu}\frac{H^2}{\phi}\left(\frac{k}{aH}\right)^{3-2\nu} \quad . \tag{55}$$

In Fig. 5 - Fig. 6 we show the behavior of the power spectrum $P_T(k,\eta)$ as a function of the wavenumber and of the parameter ω.

5.2 Spectral Index

The spectral index is obtained by the expression

$$n_T = \left.\frac{\mathrm{d}\ln P_T(k,\eta)}{\mathrm{d}\ln k}\right|_{\alpha} \quad , \tag{56}$$

where $\alpha \equiv aH = \left(\frac{2\omega+1}{2\omega-1}\right) k$ is the moment when the tensorial modes cross the horizon again. For this calculation we introduce the parameter ε, that measures the rate of decrease of the Hubble parameter during the inflation phase and it is given by

$$\varepsilon \equiv \frac{\mathrm{d}}{\mathrm{d}t}\left(\frac{1}{H}\right) = -\frac{1}{aH^2}\frac{\mathrm{d}H}{\mathrm{d}\eta} \ . \tag{57}$$

In the Brans-Dicke theory we obtain that

$$\varepsilon = \frac{2}{2\omega+1} \ . \tag{58}$$

Thus, we have

$$n_T = \frac{\mathrm{d}}{\mathrm{d}\ln k}\left[\ln\left(\frac{H^2}{\phi}\right)\right]\Bigg|_\alpha = \left(2\frac{k}{H}\frac{\mathrm{d}H}{\mathrm{d}k} - \frac{\mathrm{d}\ln\phi}{\mathrm{d}\ln k}\right)\Bigg|_\alpha \ . \tag{59}$$

The first term on the right side is calculated as follows

$$2\frac{k}{H}\frac{\mathrm{d}H}{\mathrm{d}k}\Bigg|_\alpha = 2\frac{k}{H}\frac{\mathrm{d}H}{\mathrm{d}|\eta|}\frac{\mathrm{d}|\eta|}{\mathrm{d}k}\Bigg|_\alpha = 2\frac{k}{H}\frac{\mathrm{d}H}{\mathrm{d}\eta}\left(\frac{\mathrm{d}|\eta|}{\mathrm{d}\eta}\right)^{-1}\frac{\mathrm{d}|\eta|}{\mathrm{d}k}\Bigg|_\alpha \ , \tag{60}$$

and, by using the equation (54), we have [31]

$$\frac{\mathrm{d}|\eta|}{\mathrm{d}k}\Bigg|_\alpha = \frac{\mathrm{d}}{\mathrm{d}k}\left(\left(\frac{2\omega+1}{2\omega-1}\right)\frac{1}{aH}\right)\Bigg|_\alpha = \frac{\mathrm{d}}{\mathrm{d}k}\left(\frac{1}{k}\right) = -\frac{1}{k^2} \ . \tag{61}$$

In this way, with the help of the equations (58), (60) and (61), the equation (59) can be written as

$$n_T = -2\varepsilon\left(\frac{2\omega+1}{2\omega-1}\right) - \frac{\mathrm{d}\ln\phi}{\mathrm{d}\ln k}\Bigg|_\alpha \ . \tag{62}$$

When $\omega \to \infty$ and ϕ is constant we obtain the same result that in General Relativity [31], *i. e.*, $n_T = -2\varepsilon$.

Likewise, we can calculate the term dependent on the scalar field ϕ in equation (62)

$$\frac{\mathrm{d}\ln\phi}{\mathrm{d}\ln k}\Bigg|_\alpha = \frac{k}{\phi}\frac{\mathrm{d}\phi}{\mathrm{d}k}\Bigg|_\alpha = \frac{k}{\phi}\frac{\mathrm{d}\phi}{\mathrm{d}\eta}\frac{\mathrm{d}\eta}{\mathrm{d}k}\Bigg|_\alpha = \frac{k}{\phi}\frac{\mathrm{d}\phi}{\mathrm{d}\eta}\left(\frac{\mathrm{d}|\eta|}{\mathrm{d}\eta}\right)^{-1}\frac{\mathrm{d}|\eta|}{\mathrm{d}k}\Bigg|_\alpha \ , \tag{63}$$

that with the help of equation (54) and with the background solutions of the scalar field we obtain the following result for the spectral index n_T

$$n_T = -2\varepsilon\left(\frac{2\omega+1}{2\omega-1}\right) - \frac{4}{2\omega-1} \ , \tag{64}$$

Again, the above result gives the same value of the spectral index of the tensorial modes of General Relativity at the time of inflation, when $\omega \to \infty$.

We can write the expression of n_T in terms of the parameter ω. For that we replace the relation (58) in the result found above, such that

$$n_T = \frac{-8}{2\omega - 1} \ . \tag{65}$$

By using (57) we obtain

$$2\omega = \frac{2}{\varepsilon} - 1 \ , \tag{66}$$

which can be replaced in the final result for the spectral index such that

$$n_T = \frac{4\varepsilon}{\varepsilon - 1} = -4\varepsilon(1-\varepsilon)^{-1} \ . \tag{67}$$

For small values of ε we can expand the term $(1-\varepsilon)^{-1}$ obtaining

$$n_T = -4\varepsilon(1 + \varepsilon + \mathscr{O}(\varepsilon^2)) \ , \tag{68}$$

and considering only zero-order terms we have

$$n_T \approx -4\varepsilon \ . \tag{69}$$

5.3 Energy Density

With the Lagrangian density of the gravitational waves in the Brans-Dicke theory, equation (30), we obtain the energy-momentum tensor of the gravitatinal waves

$$T_{\mu\nu} = -\frac{\phi}{8\pi}\frac{1}{4}\left(\partial_\mu h_{ij}\partial_\nu h^{ij} - \frac{1}{2}\bar{g}_{\mu\nu}\bar{g}^{\alpha\beta}\partial_\alpha h_{ij}\partial_\beta h^{ij}\right) \ . \tag{70}$$

Considering

$$h_{ij} = \sqrt{16\pi}\sum_{\lambda=\otimes,\oplus}{}^{(\lambda)}\varepsilon_{ij}h^{(\lambda)} \ , \ h^{(\lambda)} = \frac{\mu^{(\lambda)}}{a\sqrt{\phi}} \ , \tag{71}$$

we find that

$$T_{\mu\nu} = -2\phi\left(\partial_\mu h\partial_\nu h - \frac{1}{2}\bar{g}_{\mu\nu}\bar{g}^{\alpha\beta}\partial_\alpha h\partial_\beta h\right) \ , \tag{72}$$

where

$$h = h^{\otimes} = h^{\oplus} \ . \tag{73}$$

The energy density is the (00) component of the energy-momentum tensor of the gravitational waves

$$\rho = T_0{}^0 = \phi\left(\frac{h'^2}{a^2} + \bar{g}^{ij}\partial_i h\partial_j h\right) \ . \tag{74}$$

To calculate the quantum energy density we first substitute the following

$$h \to \hat{h} = \frac{\hat{\mu}}{a\sqrt{\phi}} \ , \tag{75}$$

such that the expectation value of the energy density is given by

$$\langle\rho\rangle = \langle 0|\,\rho\,|0\rangle = \frac{\phi}{a^2}\left(\langle 0|\,\hat{h}'\hat{h}'^{*}\,|0\rangle + \langle 0|\,\nabla\hat{h}\cdot\nabla\hat{h}^{*}\,|0\rangle\right) \ . \tag{76}$$

With the help of the expansions

$$\begin{aligned}\hat{\mu} &= \frac{1}{2}\int\frac{\mathrm{d}^3k}{(2\pi)^{3/2}}\left[\hat{\mu}_{\vec{k}}(\eta)e^{-i\vec{k}\cdot\vec{x}} + \hat{\mu}^{\dagger}_{\vec{k}}(\eta)e^{i\vec{k}\cdot\vec{x}}\right] \quad , \\ \hat{\pi} &= \frac{1}{2}\int\frac{\mathrm{d}^3k}{(2\pi)^{3/2}}\left[\hat{\pi}_{\vec{k}}(\eta)e^{-i\vec{k}\cdot\vec{y}} + \hat{\pi}^{\dagger}_{\vec{k}}(\eta)e^{i\vec{k}\cdot\vec{y}}\right] \ ,\end{aligned} \tag{77}$$

the solutions

$$\begin{aligned}\hat{\mu}_{\vec{k}} &= \hat{a}_{\vec{k}}(\eta_0)f_k(\eta) + \hat{a}^{\dagger}_{-\vec{k}}(\eta_0)f^{*}_k(\eta) \quad , \\ \hat{\pi}_{\vec{k}} &= \hat{a}_{\vec{k}}(\eta_0)g_k(\eta) + \hat{a}^{\dagger}_{-\vec{k}}(\eta_0)g^{*}_k(\eta) \ ,\end{aligned} \tag{78}$$

and the commutation relations

$$\left[\hat{a}_{\vec{k}}, \hat{a}^{\dagger}_{\vec{q}}\right] = \delta^{(3)}(\vec{k}-\vec{q}) \ , \tag{79}$$

we obtain

$$\langle 0|\,\hat{h}'\hat{h}'^{*}\,|0\rangle = \frac{1}{a^2\phi}\int\frac{\mathrm{d}^3k}{(2\pi)^3}\left\{|g_k|^2 + \left(\frac{\Phi'}{\Phi}\right)^2|f_k|^2 - \left(\frac{\Phi'}{\Phi}\right)(g_kf^{*}_k + g^{*}_kf_k)\right\}e^{-i\vec{k}\cdot(\vec{x}-\vec{y})} \tag{80}$$

$$\langle 0|\,\nabla\hat{h}\cdot\nabla\hat{h}^{*}\,|0\rangle = \frac{1}{a^2\phi}\int\frac{\mathrm{d}^3k}{(2\pi)^3}k^2|f_k|^2e^{-i\vec{k}\cdot(\vec{x}-\vec{y})} \ . \tag{81}$$

Thus, the expectation value of the energy density of the gravitational waves in the Brans-Dicke theory can be write as

$$\langle\rho\rangle = \frac{1}{a^4}\int\frac{\mathrm{d}^3k}{(2\pi)^3}\left[|g_k|^2 + \left(k^2 + \left(\frac{\Phi'}{\Phi}\right)^2\right)|f_k|^2 - \frac{\Phi'}{\Phi}(g_kf^{*}_k + g^{*}_kf_k)\right]e^{-i\vec{k}(\vec{x}-\vec{y})} \ . \tag{82}$$

Considering the presence of an isotropic background of relic gravitons we obtain

$$\langle\rho\rangle = \frac{1}{a^4}\int\frac{k^2\mathrm{d}k}{(2\pi)^3}\left[|g_k|^2 + \left(k^2 + \left(\frac{\Phi'}{\Phi}\right)^2\right)|f_k|^2 - \frac{\Phi'}{\Phi}(g_kf^{*}_k + g^{*}_kf_k)\right]\frac{2\sin kr}{kr} \ , \tag{83}$$

where $r = |\vec{x}-\vec{y}|$. In the limit $\vec{x}\to\vec{y}$ we have

$$\langle\rho\rangle = \frac{1}{a^4}\int\frac{k^2\mathrm{d}k}{2\pi^2}\left[|g_k|^2 + \left(k^2 + \left(\frac{\Phi'}{\Phi}\right)^2\right)|f_k|^2 - \frac{\Phi'}{\Phi}(g_kf^{*}_k + g^{*}_kf_k)\right] \ . \tag{84}$$

The energy density of the relic gravitons per logarithmic interval of the wavenumber will be given by

$$\frac{\mathrm{d}\langle\rho\rangle}{\mathrm{d}\ln k} = \frac{k^3}{2\pi^2a^4}\left[|g_k|^2 + \left(k^2 + \left(\frac{\Phi'}{\Phi}\right)^2\right)|f_k|^2 - \frac{\Phi'}{\Phi}(g_kf^{*}_k + f_kg^{*}_k)\right] \ . \tag{85}$$

The spectral energy density per logarithmic interval of the wavenumber is defined as

$$\Omega(k,\eta) = \frac{1}{\rho_c}\frac{\mathrm{d}\langle\rho\rangle}{\mathrm{d}\ln k} \ , \tag{86}$$

i. e.,

$$\Omega(k,\eta) = \frac{8\pi\phi^{-1}}{3H_0^2}\frac{k^3}{2\pi^2 a^4}\left[|g_k|^2 + \left(k^2 + \left(\frac{\Phi'}{\Phi}\right)^2\right)|f_k|^2 - \frac{\Phi'}{\Phi}\left(g_k f_k^* + f_k g_k^*\right)\right] \ , \tag{87}$$

where ρ_c is the critical density of the Universe.

For all the oscillation modes that are inside the Hubble radius (*i. e.* $k\eta >> 1$), we have $g_k \approx \mp ikf_k$. This implies that

$$\Omega(k,\eta) \approx \frac{8\pi k^5}{3\pi^2 H_0^2\phi a^4}|f_k|^2\left(1 + \frac{1}{2k^2}\left(\frac{\Phi'}{\Phi}\right)^2\right) \ , \tag{88}$$

and, rewriting in terms of the power spectrum we have

$$\Omega(k,\eta) \approx \frac{k^2}{12H_0^2 a^2}P_T\left(1 + \frac{1}{2k^2}\left(\frac{\Phi'}{\Phi}\right)^2\right) \ . \tag{89}$$

For modes outside the Hubble radius (*i. e.* $k\eta << 1$) we have $g_k \approx -\frac{\Phi'}{\Phi}f_k$. This result allow us to find that

$$\Omega(k,\eta) \approx \frac{8\pi k^5}{6\pi^2 H_0^2\phi a^4}|f_k|^2 \ , \tag{90}$$

which can also be rewritten as a function of the power spectrum P_T

$$\Omega(k,\eta) \approx \frac{k^2}{24H_0^2 a^2}P_T(k,\eta) \ . \tag{91}$$

We stress that when the modes re-enter the horizon the energy density is proportional to the power spectrum. The behavior of the spectral energy density are shown in Fig. 7 - Fig. 9.

6 Conclusions

In this work we have considered the quantum gravitational waves production during inflation in the Brans-Dicke theory. In general, we realize that, in contrast with the General Relativity, not only the scale factor $a(t)$ but also the Brans-Dicke scalar field $\phi(t)$ models the time evolution of the gravitational waves of cosmological origin.

In the initial inflationary phase of the Universe there is a natural initial vacuum state, from wich the particle occupation number can be determined. We computed the quantum particles number N_k created during extended inflation and the results indicate that the number predicted by this model approximates that predicted by General Relativity when the Brans-Dicke parameter ω is big enough, as we can see in Fig. 1 - Fig. 4.




Figure 7: The graviton energy density in the Brans-Dicke theory as a function of k, for small values of ω and fixed η.




Figure 8: The energy density of the gravitational waves in the Brans-Dicke theory as a function of k compared with the energy density in the General Relativity, for some different values of ω and fixed η.




Figure 9: The energy density of the gravitational waves in the Brans-Dicke theory as a function of $|\eta|$ compared with the energy density in the General Relativity, for some different values of ω and with fixed wavenunber k.

Afterwards we calculate three observable parameters: the quantum power spectrum P_k, the quantum spectral index n_T and the quantum energy density $\Omega(k,\eta)$ of the gravitational waves.

The result obtained for the power spectrum P_k shows that, different from the local observations and from the previous result acquired for the graviton number N_k, this model is comparable to the General Relativity one only if ω assumes small values like $0.5 < \omega < 1$. Big values of ω make the power spectrum to explode.

For the spectral index we obtain a different result. We can compare the outcome of the General Relativity for the spectral index n_T with that of the Brans-Dicke theory when $\omega >> 1$, result that agree with the one obtained for the quantum particle number N_k, and when $\omega \to \infty$ it yields a scale invariant perturbation.

In the case of the quatum energy density big values of ω make $\Omega(k,\eta)$ to explode and the General Relativity prediction is approximated when $1 < \omega < 10$, depending on the values of the wavenumber and the conformal time.

We think the main problem with the scenario of Brans-Dicke is valued at ω. In our study, the values vary significantly from the values obtained by traditional testing sites. The parameter ω assumes small or large values depending on the observable calculated. However, these drawbacks can be, in principle, circumvented if the gravitational coupling is scale-dependent, as suggested by considering quantum effects [32, 33]. In this work, we made no attempt to reconcile the values of ω deducted from local tests, with the corresponding values deduced from the analysis performed here. But, evidently, the problem must be addressed in order to have a completely realistic cosmological scenario. When gravitational waves are detected we can verify if ω has different values for different cosmological scales.

A complete study of the Brans-Dicke theory in terms of fluctuations of quantum origin also involves the calculation of scalar perturbations. This is a natural continuation of this work. The results of scalar perturbations during inflation using the Brans-Dicke theory can be compared with the results of General Relativity and with the available observational data, especially data from the cosmic microwave background radiation. We can reduce the number of the variables calculating the ratio between the tensor and the scalar power spectra $r_T = P_T/P_S$ and with this results to obtain direct information about the state of the inflationary universe. An other step to generalize our study is to do the same calculating for both scalar and tensor perturbation but in a model where the Brans-Dicke parameter ω varies with the scalar field ϕ, *i. e.*, $\omega = \omega(\phi)$. Such work is in preparation and will be available soon.

Acknowledgements: We thank CNPq (Brazil) and CAPES (Brazil) for partial financial support.

References

[1] R. Crittenden, R.L. Davis and P.J. Steinhardt, Astrophys. J. **417**, L13 (1993), [arXiv:astro-ph/9306027];

[2] R. Crittenden, J.R. Bond, R.L. Davis, G. Efstathiou and P.J. Steinhardt,Phys. Rev. Lett. **71**, 324 (1993), [arXiv:astro-ph/9303014];

[3] M. Giovannini, PMC Phys. **A** 4, 1 (2010), [arXiv:0901.3026v1];

[4] LIGO: www.ligo.caltech.edu/ ,
VIRGO: www.virgo.infn.it/ ,
MiniGrail: www.minigrail.nl/ ,
CLIO: www.icrr.u-tokyo.ac.jp ,
GEO 600: www.geo600.org/ ,
TAMA 300: tamago.mtk.nao.ac.jp/ ,
Gráviton Project: www.das.inpe.br/graviton/index.html ,
AIGO: www.gravity.uwa.edu.au/ ,
LCGT: gw.icrr.u-tokyo.ac.jp/lcgt/ ,
LISA: lisa.nasa.gov/ ;

[5] A. Riazuelo and J.P. Uzan, Phys. Rev. **D62**, 083506 (2000), [arXiv:astr-ph/0004156];

[6] A. Buonanno, M. Maggiore and C. Ungarelli, Phys. Rev. **D55**, 3330 (1997), [arXiv:gr-qc/9605072];

[7] M.P. Infante and N. Sánchez, Phys. Rev. **D61**, 083515 (2000), [arXiv:hep-th/9907185];

[8] M. Gasperini, Phys. Rev. **D56**, 4815 (1997), [arXiv:gr-qc/9704045];

[9] C. Brans and R. H. Dicke, Phys. Rev. **124**, 925 (1961);

[10] P.A.M. Dirac, Proc. Roy. Soc. (London) **A165**, 199 (1938);

[11] P. Jordan, Z. Physik **157**, 112 (1959);

[12] G.C. McVittie, M.N.R.A. Soc. **183**, 749 (1978);

[13] V. Faraoni, E. Gunzig and P. Nardone, Fund. Cosmic Phys. **20**, 12 (1999), [arXiv:gr-qc/9811047v1];

[14] D.I. Santiago and A.S. Silbergleit, Gen. Rel. Grav. **32**, 565 (2000), [arXiv:gr-qc/9904003v1];

[15] A.H. Guth, Phys. Rev. **D23**, 347 (1981);

[16] B. Bertotti, L. Iess and P. Tortora, Nature **425**, 374 (2003);

[17] C. M. Will, Living Rev. Relativity **4**, 4 (2001), [arXiv:gr-qc/0103036];

[18] D. La and P. J. Steinhardt, Phys. Rev. Lett. 62, 376 (1989);

[19] A. R. Liddle and D. Wands., Phys. Rev. **D** 45, 2665 (1992);

[20] A.G. Riess et al, Astron. J. **116**, 1009 (1998), [arXiv:astr-ph/9805201];

[21] R.R. Caldwell, R. Dave and P. J. Steinhardt, Phys. Rev. Lett. **80**, 1582-1585 (1998), [arXiv:astr-ph/9708069];

[22] C. Armendariz-Picon, V. Mukhanov and P. J. Steinhardt, Phys.Rev.Lett. **85**, 4438-4441 (2000), [arXiv:astr-ph/0004134];

[23] V. Faraoni, Phys.Rev. **D69**, 123520 (2004), [arXiv:gr-qc/0404078];

[24] N. Ogawa, Phys.Rev. **D62**, 085023 (2000), [arXiv:hep-th/0003288]; J.C. Fabris, S.V.B. Goncalves and P.E. de Souza, Gen.Rel.Grav. **34**, 53-63 (2002), [arXiv:astr-ph/0203441]; M. C. Bento, O. Bertolami and A. A. Sen, Phys.Rev. **D66**, 043507 (2002), [arXiv:gr-qc/0202064];

[25] J.C. Fabris, S.V.B. Gonçalves and R. de Sa Ribeiro, Grav. Cosmol. **12**, 49-54 (2006), [arXiv:astro-ph/0510779];

[26] J. D. Barrow, J. P. Mimoso and M. R. G. Maia, Phys. Rev. **D48**, 3630 (1993);

[27] J.P. Baptista, J.C. Fabris and S.V.B. Gonçalves, Astrophysics and Space Science **246**, 315-331 (1996), [arXiv:gr-qc/9603015];

[28] N. D. Birrel and P. C. W. Davies, *Quantum fields in curved space*, Cambridge University Press, Cambridge (1982);

[29] D. Baumann, *TASI Lectures on Inflation*, Lectures da 2009 Theoretical Advanced Study Institute na Univ. of Colorado, Boulder, [arXiv:0907.5424v1];

[30] M. Abramowitz e I. A. Stegun, *Handbook of Mathematical Functions*, National Bureau of Standards, (1964);

[31] S. Dodelson, *Modern Cosmology*, Academic Press, (2003);

[32] M. Reuter and H. Weyer, Phys. Rev. **D70**, 124028 (2004), [arXiv:hep-th/0410117];

[33] I.L. Shapiro, J. Sola and H. Stefancic, JCAP **0501**, 012 (2005), [arXiv:hep-ph/0410095].